

Experts in human cell culture

Human primary cells, stem cells, blood and immune cells, and optimized cell culture media



Table of content

| | |
|--|----|
| Why choose us? | 05 |
| Our locations worldwide | 05 |
| We understand your cell culture needs | 06 |
| Excipient GMP-grade media | 06 |
| Customized media according to your needs | 06 |
| Customized cell culture | 07 |
| Scientific advice and support | 07 |
| Our portfolio of human cell cultures | 09 |
| Human primary cells | 09 |
| Cell culture media | 09 |
| <i>In vitro</i> 3D cell models | 10 |
| Cancer cell culture | 11 |
| Our variety of human primary cells | 13 |
| Our mesenchymal stem cells | 14 |
| MSC sources | 15 |
| MSC culture | 15 |
| Our variety of blood and immune cell types | 16 |
| Cell types and media overview | 17 |
| Accelerate your cell and gene therapy development | 19 |
| Discover our customization possibilities | 19 |
| Our products | 21 |
| Human primary cell culture products | 21 |
| Human stem cell culture products | 25 |
| Human blood and immune cell culture products | 26 |
| Cancer cell culture products | 27 |
| Reagents for cell dissociation and cryopreservation | 28 |
| References | 30 |



Why choose us?

We empower scientists to conduct world-class research by offering an extensive portfolio of human primary cells, stem cells, blood cells, and optimized cell growth and differentiation media. We can fine-tune our flexible manufacturing process to offer customized products that meet your research and manufacturing needs, regardless of the organization's size and stage of your research. For over 30 years, we have continually supplied scientists with the tools and support they need to conduct groundbreaking research.

Our commitment to quality

1

Our **trained scientists** are here to support you at every step of your research.

2

Our products meet European and international **quality and ethical standards**.

3

We own the entire manufacturing and tissue procurement process, which allows us to offer **custom cells and media**.

4

We have **35 years of expertise** and offer products that are used in over 1,000 peer-reviewed articles every year.

5

Our **ISO 9001:2015 certification** ensures that we provide products that meet your needs and regulatory requirements.

6


Our quality management system is certified according to the **EXCiPACT™ Good Manufacturing Practice (GMP) standard**.

7

Our wealth of resources ensures **easy access** to information related to our products.

Our locations worldwide

We operate in 38 countries across three continents. This enables us to provide you with all the support you need, quickly and reliably.



- | | | | | | | | |
|----------|----------------|---------|---------------|-------------|-------------|-------------|----------------|
| Austria | Czech Republic | Greece | Italy | Monaco | Romania | Spain | United Kingdom |
| Belgium | Denmark | Hungary | Japan | Netherlands | Singapore | Sweden | United States |
| Bulgaria | Finland | India | Liechtenstein | Norway | Slovakia | Switzerland | |
| Canada | France | Ireland | Luxembourg | Poland | Slovenia | Taiwan | |
| Croatia | Germany | Israel | Malaysia | Portugal | South Korea | Thailand | |

We understand your cell culture needs

We're here to support every stage of your cell culture workflow, from research to manufacturing.

Excipient GMP-grade media

Our quality management system is certified according to EXCiPACT™, an internationally recognized GMP standard. Our entire product development pipeline is compliant with international regulatory guidelines.

This means that we can offer media and reagents produced in a controlled environment that meets your regulatory requirements. Contact us to discuss your needs and regulatory requirements.



- Our quality management system is certified according the EXCiPACT™ GMP standard




- PromoCell operates according to ISO 9001:2015 in order to consistently provide products and services that meet customer requirements as well as applicable statutory and regulatory requirements.

Customized media according to your needs

The commercial and clinical translation of cell-based therapies requires individualized solutions that span small-scale non-GMP development to GMP compliant scaled-up

manufacturing. Our flexible manufacturing pipeline enables us to offer a wide range of customized cell culture media.



Customized cell culture

In addition to our extensive stock of cells from matched donors, we can also isolate cells from specific donors who match your donor specifications.

Do you need cells with a specific HLA type for cytotoxicity screening? Contact us to discuss our large selection of cell types from HLA-typed donors.

We also offer a standardized cell expansion service if you need a large number of cells. After we expand the cells using our standardized expansion protocols, we perform an additional quality control so that you can start your experiments right away.

Scientific advice and support

Our qualified and highly trained cell culture experts can help you elevate your cell culture research by providing scientific support at every step.

Do you need training to handle valuable primary cells? Do you struggle with cell contamination? Contact us to discuss how we can address your cell culture concerns.





Our portfolio of human cell cultures


Human primary cells

We provide primary cells from various tissues and organs. All our cells are directly isolated from tissues and retain the morphological and functional characteristics of their tissue of origin.

In addition to human primary cells from a wide variety of organs and tissues, we also offer human stem, blood, and immune cells from the bone marrow, umbilical cord, placenta, adipose tissue, peripheral blood, and cord blood. To support you with the culture of human cells,


we also offer high-quality media that simulate the *in vivo* environment or promote cell differentiation.

We manufacture all our primary cells and stem cells according to the highest ethical and legal standards. When obtaining human tissues from donors, we follow strict European biomedical conventions, ethical standards, and privacy regulations.



Physiologically relevant solutions

- Human primary cells feature tissue characteristics close to *in vivo* conditions.



Individual donor characteristics

- For studying cells from donors of different genders, ages, and ethnicities.



Verified cell identity and function

- Our strict quality control ensures correct cellular identity, growth, and differentiation performance.

Cell culture media

Using high-quality media is vital for successful cell culture. To ensure that we provide you with high-quality media, we adhere to strict quality standards and produce media using only raw materials from strictly approved sources.

We are committed to providing media that is standardized and optimized for your needs and have replaced serum with well-defined substances wherever possible. Using serum-free media minimizes the risk of introducing adventitious agents to your cell cultures, provides lot-to-lot consistency, and

simplifies regulatory submissions. The use of our serum-free media also ensures optimized cell growth while avoiding the unpredictable effects of indeterminate compounds.


Time is your most valuable asset. Setting up a standardized and robust cell culture or differentiation protocol can be laborious and time-consuming. To help you focus on your research rather than optimizing cell differentiation protocols, we offer ready-to-use differentiation media, along with detailed cell culture protocols and application notes.

In vitro **3D cell models**


Generating physiologically relevant data relies on the use of models that closely resemble the biological systems of interest. 3D cell models allow for a better understanding of complex biology in a physiologically relevant context. We have the expertise and appropriate systems to support your journey

toward the future of *in vitro* research.


From spheroids to organoids, air-liquid interface culture, and magnetic 3D culture, we offer all the reagents, detailed cell culture protocols, and application notes to help you establish your 3D cell models.



Spheroids



Organoids



Air-liquid interface




Magnetic 3D cell culture



Cancer cell culture


We offer a portfolio of cancer cell media to help you establish primary cancer cell lines from biopsies, develop 3D tumor models, or build screening assays.



Our **Primary Cancer Culture System**, **3D Tumorsphere Medium XF**, and **Cancer Cell Line Medium XF** are ideal for culturing cancer stem cells (CSCs). Despite making up 1–2% of all cells in a tumor, CSCs play a critical role in tumor development, progression, and metastasis. Studying CSCs is, therefore,

essential for understanding cancer biology and for developing effective therapies. All three media types are synthetic, animal component-free, and xeno-free while containing all the nutrients necessary for the growth and long-term cultivation of CSCs in 2D or 3D cultures. You can also use these

media to culture most established cell lines and to establish cell lines from biopsies of various cancer types. In addition to CSC research, our cancer cell culture media can be used for drug discovery and cell-based therapy development using co-cultures of cancer and immune cells.



Our variety of human primary cells

Established cell lines are an important tool for biomedical research but do not necessarily reflect the biology and behavior of cells in tissues and organs. In contrast, primary cells closely resemble the genetic makeup and behavior of the tissue of origin. This makes human primary cells a physiologically relevant cell model for studying complex biological processes and human diseases.

The process of isolating and culturing

cells from tissues and organs is complex and can introduce variations in your data. To ensure that your research is reproducible, our experts use highly standardized reagents and protocols when isolating primary cells from donors.

We offer a wide variety of high-quality human primary cells. We isolate our primary cells from tissues obtained from approved medical centers that comply with strict ethical

and regulatory standards. All our primary cells are available as either cryopreserved or proliferating cells. Each cell batch is tested for cell type-specific markers, morphology, population doubling time, and proliferation capacity. Additionally, we test all our primary cells to confirm that they are not infected with HIV1, HIV2, HBV, HCV, fungi, or mycoplasma.

| | | | | |
|---|---|---|---|---|
|  <p>Musculoskeletal system</p> <p>Skeletal Muscle Cells Chondrocytes Osteoblasts</p> |  <p>Skin and connective tissue</p> <p>Fibroblasts Keratinocytes Melanocytes Dermal Endothelial Cells Follicle Dermal Papilla Cells</p> |  <p>Uterus</p> <p>Smooth Muscle Cells Fibroblasts Endothelial Cells</p> |  <p>Bone marrow</p> <p>Mesenchymal Stem Cells</p> |  <p>Kidney</p> <p>Renal Epithelial Cells</p> |
|  <p>Cardiovascular system</p> <p>Smooth Muscle Cells Endothelial Cells Myocytes Fibroblasts</p> |  <p>Umbilical cord and placenta</p> <p>Pericytes Mesenchymal Stem Cells Mononuclear Cells Smooth Muscle Cells CD34⁺ Progenitor Cells</p> |  <p>Respiratory system</p> <p>Airway Epithelial Cells Fibroblasts Endothelial Cells Smooth Muscle Cells</p> |  <p>Adipose tissue</p> <p>Preadipocytes Mesenchymal Stem Cells</p> |  <p>Peripheral blood</p> <p>Macrophages Mononuclear Cells</p> |

HLA-typed cells

All our human primary cells are sourced from HLA-typed donors. Along with the cells, we also offer HLA Typing Certificate with high-resolution (4-digit) typing results for HLA-A, -B, -C, -DPA1/DPB1, -DQA1/DQB1, and -DRB1/DRB345.

Primary cells from donors with a known disease

We offer a broad range of cells from donors who have the disease of your research focus. These cells feature the same specifications and undergo the same rigid quality control tests as our primary cells from normal donors. Our quality control tests include marker characterization, growth performance, and morphological characterization.

Our mesenchymal stem cells

We offer a collection of human mesenchymal stem cells isolated from the bone marrow, umbilical cord, and adipose tissue. All cells are obtained from approved donor programs at associated medical centers. We also offer optimized differentiation media and high-quality media that simulate the *in vivo* environment of primary cells.



Bone marrow

Human Mesenchymal Stem Cells from Bone Marrow (hMSC-BM)

Cat. No. C-12974
Marker expression:
CD105⁺ | CD73⁺ | CD90⁺
CD45⁻ | CD34⁻ |
CD14⁻ | CD19⁻ | HLA-DR⁻



Adipose tissue

Human Mesenchymal Stem Cells from Adipose Tissue (hMSC-AT)

Cat. No. C-12977
Marker expression:
CD105⁺ | CD73⁺ | CD90⁺
CD45⁻ | CD34⁻ |
CD14⁻ | CD19⁻ | HLA-DR⁻



Umbilical cord

Human Mesenchymal Stem Cells from Umbilical Cord Matrix (hMSC-UC)

Cat. No. C-12971
Marker expression:
CD105⁺ | CD73⁺ | CD90⁺
CD45⁻ | CD34⁻ |
CD14⁻ | CD19⁻ | HLA-DR⁻


Key advantages of our mesenchymal stem cells

- **Verified expression of surface markers** that meet the International Society for Cellular Therapy (ISCT) criteria
- **Standardized and well-characterized**, enabling reproducible research
- Possibility of **large-scale** production
- **Confirmed differentiation** into osteoblasts, adipocytes, and chondrocytes
- Expression of **trilineage surface markers** when cultured in our optimized media
- **Custom media** according to your needs and regulatory requirements
- **Excipient GMP-grade media**

MSC sources

Stem cells are often poorly characterized, and their use requires time-consuming and standardized protocols. To support translation of cellular therapies, the International Society for Cell and Gene Therapy (ISCT) has described three key characteristics of MSCs:

1 Adherence to plastic




MSCs must adhere to plastic to grow under standard culture conditions

2 Marker expression

| Positive ≥95% + | Negative ≤2% + |
|----------------------|---|
| ■ CD105 ⁺ | ■ CD45 ⁻ |
| ■ CD73 ⁺ | ■ CD34 ⁻ |
| ■ CD90 ⁺ | ■ CD14 ⁻ or CD11b ⁻ |
| | ■ CD79α ⁻ or CD19 ⁻ |
| | ■ HLA-DR ⁻ |

MSCs are characterized by the presence or absence of specific surface markers

3 Multipotent differentiation potential



MSCs are multipotent and can differentiate into various cell lineages

MSC culture

Our optimized growth media ensure consistent growth and maintenance of the trilineage differentiation potential. Our media contain all the growth factors and supplements necessary for the robust expansion of human mesenchymal stem cells and the generation of high-quality multipotent cells.




Fig. 3: In vitro differentiation of hMSCs into adipocytes, chondrocytes, osteoblasts, and neuronal cells. **A:** Accumulation of lipid vesicles in adipocytes differentiated from bone marrow-derived hMSCs. The cells show typical characteristics of mature adipocytes, including intracellular lipid vacuoles. **B:** Alcian blue staining of cartilage spheroids after in vitro differentiation of hMSCs. Alcian blue stains cartilage extracellular matrix. **C:** Alizarin Red S staining of mature osteoblasts after in vitro differentiation of bone marrow-derived hMSCs. Alizarin Red S stains extracellular calcium deposits. **D:** Representative image of neurons after in vitro differentiation of bone marrow-derived hMSCs into neurogenic lineages.

Our variety of blood and immune cell types

The immune system is complex, and studying immunological and hematologic diseases can be challenging. Immune cells, such as macrophages and dendritic cells, are a valuable model for the characterization of the mechanisms contributing to the development of autoimmune diseases, inflammation, and allergies. Immune cells are also ideal for the development of novel therapies to inhibit

inflammation, fight infections, or overcome immunosuppression.

Co-cultures of different immune cells and cancer cells can also be used to study the complex interactions between cancer cells and immune cells, as well as to identify new ways to enhance cancer cell killing by immune cells.

CD34⁺ hematopoietic stem cells are an

excellent source of erythrocytes, macrophages, dendritic cells, and lymphocytes. Immune cells can also be isolated from human peripheral blood mononuclear cells (PBMCs). To elevate your immunology research, we offer a wide range of progenitor cells and differentiated blood cells, along with optimized media that mimic their *in vivo* environment or stimulate cell differentiation.



Peripheral blood cells

Human Mononuclear Cells from Peripheral Blood (hMNC-PB)

Cat. No. C-12907
Marker expression:
CD31⁻ | CD34⁻
CD105⁺ | CD146⁺

Human M1- and M2 Macrophages (GM-CSF/M-CSF)

Cat. No. C-12914/C-12915
Marker expression:
M1: CD80⁺ | CD68⁺
M2: CD163⁺ | CD68⁺

Human CD14⁺ Monocytes from Peripheral Blood (hMoCD14⁺-PB)

Cat. No. C-12909
Marker expression:
CD14⁺



Umbilical cord blood cells

Human Mononuclear cells from Umbilical Cord Blood (hMNC-CB)







Cat. No. C-12901
Marker expression:
CD31⁻ | CD34⁻
CD105⁺ | CD146⁺

Human CD34⁺ Progenitor Cells from Umbilical Cord Blood (hCD34⁺-CB)

Cat. No. C-12921
Marker expression:
CD34⁺



Cell types and media overview

| Cells  | | Blood and immune cells | Expansion media | Differentiation media |
|---|--|------------------------|-----------------|-----------------------|
|  | Macrophage generation (M1 / M2) Straightforward and efficient differentiation of highly pure M1- or M2-macrophages directly from freshly isolated peripheral blood monocytes. | ✓ | | ✓ |
|  | Dendritic cell generation Easy and efficient generation of immature and fully mature myeloid dendritic cells from peripheral blood monocytes. | | | ✓ |
|  | CD34⁺ hematopoietic progenitor cell culture Isolated by positive immunomagnetic selection from fresh cord blood. Suitable xeno-free HPC expansion medium is also available. | ✓ | ✓ | |
|  | Mononuclear cell culture Mononuclear cells (MNC) represent the lymphocyte and monocyte fraction isolated from fresh umbilical cord blood or peripheral blood. | ✓ | ✓ [*] | |
|  | Monocyte cell culture CD14 ⁺ monocytes are isolated from fresh peripheral blood by positive selection with over 95% purity. | ✓ | ✓ [*] | |

*short-term maintenance medium for up to 48 hours





Accelerate your cell and gene therapy development

Cell and gene therapies are becoming promising treatment options for several diseases. However, cell and gene therapies are more complex than conventional drugs, and their efficacy and safety may vary depending on many factors.

The use of standardized and well-characterized primary cells and optimized cell culture

media is key to the success of cell and gene therapies and a requirement by regulatory bodies.


We can help you accelerate your cell and gene therapy development by offering a broad range of standardized cell cultures and customized media. Moreover, we provide scientific and regulatory support at every step

in cell and gene therapy development.

Contact us to help you find the cell culture solution that fits your needs and is optimized for your application:

info@promocell.com

Discover our customization possibilities





Our products

Human primary cell culture products

Primary cells

| Cell type | Description | Catalog no. | Marker | Recommended culture media |
|-----------------------------------|--|-------------|--|---------------------------|
| Cardiac myocytes | Human Cardiac Myocytes (HCM) | C-12810 | GATA-4 ⁺ and Sarcomeric α -actin ⁺ | C-22070 |
| Chondrocytes | Human Chondrocytes (HCH) | C-12710 | Differentiation into 3D spheroids tested | C-27101 |
| Endothelial cells (large vessels) | Human Umbilical Vein Endothelial Cells (HUVEC) single donor | C-12200 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ | C-22010 C-22011 |
| | Human Umbilical Vein Endothelial Cells (HUVEC) pooled | C-12203 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ | C-22010 C-22011 |
| | Human Umbilical Vein Endothelial Cells (HUVEC) isolated in Growth Medium 2, single donor | C-12206 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ | C-22011 |
| | Human Umbilical Vein Endothelial Cells (HUVEC) isolated in Growth Medium 2, pooled | C-12208 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ | C-22011 |
| | Human Umbilical Vein Endothelial Cells (HUVEC) pre-screened | C-12205 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ VEGF response | C-22010 C-22011 |
| | Human Umbilical Artery Endothelial Cells (HUAEC) | C-12202 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ | C-22010 C-22011 |
| | Human Aortic Endothelial Cells (HAoEC) | C-12271 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ | C-22020 C-22022 |
| | Human Coronary Artery Endothelial Cells (HCAEC) | C-12221 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ | C-22020 C-22022 |
| | Human Pulmonary Artery Endothelial Cells (HPAEC) | C-12241 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ | C-22010 C-22011 |
| | Human Saphenous Vein Endothelial Cells (HSAVEC) | C-12231 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ | C-22010 C-22011 |
| Endothelial cells (microvascular) | Human Dermal Microvascular Endothelial Cells (HDMEC) juvenile foreskin | C-12210 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ | C-22020 C-22022 |
| | Human Dermal Microvascular Endothelial Cells (HDMEC) adult donor | C-12212 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ | C-22020 C-22022 |
| | Human Dermal Microvascular Endothelial Cells (HDMEC) pre-screened | C-12215 | CD31 ⁺ DiI-Ac-LDL uptake ⁺ VEGF response | C-22020 C-22022 |
| | Human Dermal Blood Endothelial Cells (HDBEC) juvenile foreskin | C-12211 | Podoplanin ⁻ CD31 ⁺ | C-22020 |
| | Human Dermal Blood Endothelial Cells (HDBEC) adult donor | C-12225 | Podoplanin ⁻ CD31 ⁺ | C-22020 |

| Cell type | Description | Catalog no. | Marker | Recommended culture media |
|-----------------------------------|---|-------------|---|---------------------------|
| Endothelial cells (microvascular) | Human Dermal Lymphatic Endothelial Cells (HDLEC) juvenile foreskin | C-12216 | Podoplanin ⁺ CD31 ⁺ | C-22022 |
| | Human Dermal Lymphatic Endothelial Cells (HDLEC) adult donor | C-12217 | Podoplanin ⁺ CD31 ⁺ | C-22022 |
| | Human Cardiac Microvascular Endothelial Cells (HCMEC) | C-12285 | CD31 ⁺ Dil-Ac-LDL uptake ⁺ | C-22020 C-22022 |
| | Human Pulmonary Microvascular Endothelial Cells (HPMEC) | C-12281 | CD31 ⁺ Dil-Ac-LDL uptake ⁺ | C-22020 C-22022 |
| | Human Uterine Microvascular Endothelial Cells (HUtMEC) | C-12295 | CD31 ⁺ Dil-Ac-LDL uptake ⁺ | C-22020 C-22022 |
| Epithelial cells | Human Nasal Epithelial Cells (HNEpC) | C-12620 | Cytokeratin ⁺ | C-21060 |
| | Human Tracheal Epithelial Cells (HTEpC) | C-12644 | Cytokeratin ⁺ | C-21060 |
| | Human Bronchial Epithelial Cells (HBEpC) | C-12640 | Cytokeratin ⁺ | C-21060 |
| | Human Small Airway Epithelial Cells (HSAEpC) | C-12642 | Cytokeratin ⁺ | C-21070 |
| | Human Renal Epithelial Cells (HREpC) | C-12665 | Cytokeratin ⁺ | C-26030 |
| | Human Renal Cortical Epithelial Cells (HRCEpC) | C-12660 | Cytokeratin ⁺ | C-26030 |
| Fibroblasts | Normal Human Dermal Fibroblasts (NHDF) juvenile foreskin | C-12300 | CD90 ⁺ | C-23010 |
| | Normal Human Dermal Fibroblasts (NHDF) adult donor | C-12302 | CD90 ⁺ | C-23020 |
| | Human Pulmonary Fibroblasts (HPF) | C-12360 | CD90 ⁺ | C-23020 |
| | Human Aortic Adventitial Fibroblasts (HAoAF) | C-12380 | CD90 ⁺ | C-23020 |
| | Human Cardiac Fibroblasts (HCF) | C-12375 | CD90 ⁺ Fibronectin ⁺ | C-23025 |
| | Human Uterine Fibroblasts (HUF) | C-12385 | CD90 ⁺ | C-23020 |
| Follicle dermal papilla cells | Human Follicle Dermal Papilla Cells (HFDPC) | C-12071 | Alkaline phosphatase ⁺ | C-26501 |
| Keratinocytes | Normal Human Epidermal Keratinocytes (NHEK) juvenile foreskin, single donor | C-12001 | Cytokeratin ⁺ | C-20011 |
| | Normal Human Epidermal Keratinocytes (NHEK) juvenile foreskin, pooled | C-12005 | Cytokeratin ⁺ | C-20011 |
| | Normal Human Epidermal Keratinocytes (NHEK) adult, single donor | C-12003 | Cytokeratin ⁺ | C-20011 |

| Cell type | Description | Catalog no. | Marker | Recommended culture media |
|-----------------------|---|-------------|---|-------------------------------|
| Keratinocytes | Normal Human Epidermal Keratinocytes (NHEK) adult, pooled | C-12006 | Cytokeratin ⁺ | C-20011 |
| | Normal Human Epidermal Keratinocytes (NHEK) GM3 juvenile foreskin, single donor | C-12011 | Cytokeratin ⁺ | C-20021 |
| | Normal Human Epidermal Keratinocytes (NHEK) GM3 juvenile foreskin, pooled | C-12015 | Cytokeratin ⁺ | C-20021 |
| | Normal Human Epidermal Keratinocytes (NHEK) GM3 adult, single donor | C-12013 | Cytokeratin ⁺ | C-20021 |
| | Normal Human Epidermal Keratinocytes (NHEK) GM3 adult, pooled | C-12016 | Cytokeratin ⁺ | C-20021 |
| Melanocytes | Normal Human Epidermal Melanocytes (NHEM) juvenile foreskin | C-12400 | Mel-5 ⁺ | C-24010 |
| | Normal Human Epidermal Melanocytes (NHEM) juvenile foreskin cultured in M3 Medium | C-12422 | Mel-5 ⁺ | C-24310 |
| | Normal Human Epidermal Melanocytes (NHEM) adult donor, cultured in M3 Medium | C-12413 | Mel-5 ⁺ | C-24310 |
| Osteoblasts | Human Osteoblasts (HOB) | C-12720 | Alkaline phosphatase ⁺ Mineralization tested | C-27001 C-27020 |
| Preadipocytes | Human White Preadipocytes (HWP) subcutaneous | C-12735 | Differentiation tested | C-27410 C-27436 C-27438 |
| | Human White Preadipocytes (HWP) visceral | C-12732 | Differentiation tested | C-27410 C-27436 C-27438 |
| Skeletal muscle cells | Human Skeletal Muscle Cells (SkMC) | C-12530 | Differentiation capacity to multinucleate syncytia tested | C-23060 C-23061 |
| Smooth muscle cells | Human Aortic Smooth Muscle Cells (HAoSMC) | C-12533 | Smooth muscle α -actin ⁺ CD31 ⁻ | C-22062 |
| | Human Coronary Artery Smooth Muscle Cells (HCASMC) | C-12511 | Smooth muscle α -actin ⁺ CD31 ⁻ | C-22062 |
| | Human Pulmonary Artery Smooth Muscle Cells (HPASMC) | C-12521 | Smooth muscle α -actin ⁺ CD31 ⁻ | C-22062 |
| | Human Umbilical Artery Smooth Muscle Cells (HUASMC) | C-12500 | Smooth muscle α -actin ⁺ CD31 ⁻ | C-22062 |
| | Human Tracheal Smooth Muscle Cells (HTSMC) | C-12565 | Smooth muscle α -actin ⁺ CD31 ⁻ | C-22062 |
| | Human Bronchial Smooth Muscle Cells (HBSMC) | C-12561 | Smooth muscle α -actin ⁺ CD31 ⁻ | C-22062 |
| | Human Uterine Smooth Muscle Cells (HuTSMC) | C-12575 | Smooth muscle α -actin ⁺ CD31 ⁻ | C-22062 |

Cell culture media for expansion and differentiation

| Cell type | Product | Size (mL) | Catalog no. |
|--------------------------------------|---|-----------|------------------------------------|
| Cardiac myocytes | Myocyte Growth Medium | 500 | Ready-to-use C-22070 / Kit C-22170 |
| Chondrocytes | Chondrocyte Growth Medium | 500 | C-27101 |
| Endothelial cells (large vessels) | Endothelial Cell Growth Medium | 500 | Ready-to-use C-22010 / Kit C-22110 |
| | Endothelial Cell Growth Medium 2 | 500 | Ready-to-use C-22011 / Kit C-22111 |
| Endothelial cells (microvascular) | Endothelial Cell Growth Medium MV | 500 | Ready-to-use C-22020 / Kit C-22120 |
| | Endothelial Cell Growth Medium MV 2 | 500 | Ready-to-use C-22022 / Kit C-22121 |
| Epithelial cells | Airway Epithelial Cell Growth Medium | 500 | Ready-to-use C-21060 / Kit C-21160 |
| | Small Airway Epithelial Cell Growth Medium | 500 | Ready-to-use C-21070 / Kit C-21170 |
| | Air-Liquid Interface Medium (ALI-Airway) | 500 | C-21080 |
| | Renal Epithelial Cell Growth Medium 2 | 500 | Ready-to-use C-26030 / Kit C-26130 |
| Fibroblasts | Fibroblast Growth Medium | 500 | Ready-to-use C-23010 / Kit C-23110 |
| | Fibroblast Growth Medium 2 | 500 | Ready-to-use C-23020 / Kit C-23120 |
| | Fibroblast Growth Medium 3 | 500 | Ready-to-use C-23025 / Kit C-23130 |
| | Fibroblast Growth Medium D-ACF | 500 | C-23027 |
| | Fibroblast Growth Medium D-ACF, phenol red-free | 500 | C-23017 |
| Follicle dermal papilla cells | Follicle Dermal Papilla Cell Growth Medium | 500 | Ready-to-use C-26501 / Kit C-26502 |
| Keratinocytes | Keratinocyte Growth Medium 2 | 500 | Ready-to-use C-20011 / Kit C-20111 |
| | Keratinocyte Growth Medium 3 | 500 | C-20021 |
| Melanocytes | Melanocyte Growth Medium | 500 | Ready-to-use C-24010 / Kit C-24110 |
| | Melanocyte Growth Medium M3 | 500 | C-24310 |
| Osteoblasts | Osteoblast Growth Medium | 500 | C-27001 |
| | Osteoblast Mineralization Medium | 500 | C-27020 |

| Cell type | Product | Size (mL) | Catalog no. |
|-----------------------|---|-----------|------------------------------------|
| Preadipocytes | Preadipocyte Growth Medium | 500 | Ready-to-use C-27410 / Kit C-27417 |
| | Preadipocyte Differentiation Medium | 500 | Ready-to-use C-27436 / Kit C-27437 |
| | Adipocyte Nutrition Medium | 500 | Ready-to-use C-27438 / Kit C-27439 |
| Skeletal muscle cells | Skeletal Muscle Cell Growth Medium | 500 | Ready-to-use C-23060 / Kit C-23160 |
| | Skeletal Muscle Cell Differentiation Medium | 500 | C-23061 |
| Smooth muscle cells | Smooth Muscle Cell Growth Medium 2 | 500 | Ready-to-use C-22062 / Kit C-22162 |

Human stem cell culture products

Human stem and progenitor cells

| Cell type | Description | Catalog no. | Marker | Recommended culture media |
|--------------------------------|---|-------------|--|---|
| Human mesenchymal stem cells | Human Mesenchymal Stem Cells from Bone Marrow (hMSC-BM) | C-12974 | CD105 ⁺ /CD73 ⁺ /CD90 ⁺ and CD45 ⁻ /CD34 ⁻ /CD14 ⁻ /CD19 ⁻ /HLA-DR ⁻ | C-28009 C-28019 EQ-C-28019 C-28018 |
| | Human Mesenchymal Stem Cells from Umbilical Cord Matrix (hMSC-UC) | C-12971 | CD105 ⁺ /CD73 ⁺ /CD90 ⁺ and CD45 ⁻ /CD34 ⁻ /CD14 ⁻ /CD19 ⁻ /HLA-DR ⁻ | C-28009 C-28019 EQ-C-28019 C-28018 |
| | Human Mesenchymal Stem Cells from Adipose Tissue (hMSC-AT) | C-12977 | CD105 ⁺ /CD73 ⁺ /CD90 ⁺ and CD45 ⁻ /CD34 ⁻ /CD14 ⁻ /CD19 ⁻ /HLA-DR ⁻ | C-28009 C-28019 EQ-C-28019 C-28018 |
| Hematopoietic progenitor cells | Human CD34 ⁺ Progenitor Cells from Cord Blood (hCD34 ⁺ -CB) | C-12921 | CD34 ⁺ | C-28021 C-39891 |
| Pericytes | Human Pericytes from Placenta (hPC-PL) | C-12980 | CD105 ⁺ /CD146 ⁺ and CD31 ⁻ /CD34 ⁻ | C-28041 |

Cell culture media for expansion and differentiation

| Cell type | Product | Size (mL) | Catalog no. |
|--------------------------------|---|-----------|-------------|
| Human mesenchymal stem cells | PromoExQ_MSC Growth Medium XF | 500 | EQ-C-28019 |
| | Mesenchymal Stem Cell Growth Medium 2 (Ready-to-use) | 500 | C-28009 |
| | Mesenchymal Stem Cell Growth Medium 2, phenol red-free | 500 | C-28017 |
| | Mesenchymal Stem Cell Growth Medium XF (Ready-to-use) | 500 | C-28019 |
| | Mesenchymal Stem Cell Growth Medium XF (Ready-to-use), phenol red-free | 500 | C-28018 |
| | Mesenchymal Stem Cell Adipogenic Differentiation Medium 2 (Ready-to-use) | 100 | C-28016 |
| | Mesenchymal Stem Cell Chondrogenic Differentiation Medium (Ready-to-use) | 100 | C-28012 |
| | Mesenchymal Stem Cell Osteogenic Differentiation Medium (Ready-to-use) | 100 | C-28013 |
| | Mesenchymal Stem Cell Chondrogenic Differentiation Medium w/o Inducers (Ready-to-use) | 100 | C-28014 |
| | Mesenchymal Stem Cell Neurogenic Differentiation Medium (Ready-to-use) | 100 | C-28015 |
| Hematopoietic progenitor cells | Hematopoietic Progenitor Expansion Medium XF | 500 | C-28021 |
| | Hematopoietic Progenitor Cell Expansion Medium XF, phenol red-free | 500 | C-28022 |
| | Cytokine Mix E for 100 ml HPC Expansion Medium XF | 1 | C-39890 |
| | Cytokine Mix E for 500 ml HPC Expansion Medium XF | 5 | C-39891 |
| Pericytes | Pericyte Growth Medium 2 (Ready-to-use) | 500 | C-28041 |
| | Pericyte Growth Medium 2, phenol red-free | 500 | C-28042 |

Human blood and immune cell culture products

Human blood and immune cells

| Cell type | Description | Catalog no. | Marker | Recommended culture media |
|-------------|--|-------------|---|---------------------------|
| Macrophages | Human M1 Macrophages (GM-CSF), monocyte-derived, single donor (hMDM-GMCSF(-)) | C-12914 | CD80 ⁺ CD68 ⁺ | C-28055 |
| | Human M2 Macrophages (M-CSF), monocyte-derived, single donor (hMDM-MCSF(-)) | C-12915 | CD163 ⁺ CD68 ⁺ | C-28056 |
| Monocytes | Human CD14 ⁺ Monocytes from Peripheral Blood (hMoCD14 ⁺ -PB), single donor | C-12909 | CD14 ⁺ | C-28030 |

| Cell type | Description | Catalog no. | Marker | Recommended culture media |
|-------------------|---|-------------|--|---------------------------|
| Mononuclear cells | Human Mononuclear Cells from Peripheral Blood (hMNC-PB), single donor, ultra-pure | C-12907 | Lymphocytes Monocytes Granulocytes | C-28030 |
| | Human Mononuclear Cells from Cord Blood (hMNC-CB), single donor, ultra-pure | C-12901 | Lymphocytes Monocytes Granulocytes | C-28030 |

Cell culture media for expansion and differentiation

| Cell type | Product | Size (mL) | Catalog no. |
|-------------------|------------------------------------|-----------|-------------|
| Dendritic cells | DC Generation Medium | 250 | C-28050 |
| | DC Generation Medium XF | 250 | C-28052 |
| Macrophages | M1-Macrophage Generation Medium XF | 250 | C-28055 |
| | M2-Macrophage Generation Medium XF | 250 | C-28056 |
| Monocytes | Monocyte Attachment Medium | 250 | C-28051 |
| Mononuclear cells | Mononuclear Cell Medium | 500 | C-28030 |

Cancer cell culture products

Cancer cell culture media

| Product | Size (mL) | Catalog no. |
|--|-----------|-------------|
| Primary Cancer Culture System | 250 | C-28081 |
| NCCD Reagent | 2 | C-43080 |
| 3D Tumorsphere Medium XF | 250 | C-28070 |
| 3D Tumorsphere Medium XF (Ready-to-use), phenol red-free | 250 | C-28075 |
| Cancer Cell Line Medium XF | 250 | C-28077 |

Reagents for cell dissociation and cryopreservation

| Product | Size (mL) | Catalog no. |
|--------------------------------|-----------|-------------|
| DetachKit | 30 | C-41200 |
| | 125 | C-41210 |
| | 250 | C-41220 |
| Macrophage Detachment Solution | 250 | C-41330 |
| Accutase-Solution | 100 | C-41310 |
| Cryo-SFM Plus | 30 | C-29920 |
| | 125 | C-29922 |
| Cryo-SFM Plus, phenol red-free | 30 | C-29930 |
| | 125 | C-29932 |



References

1. Ahrlund-Richter L, De Luca M, Marshak DR, Munsie M, Veiga A, Rao M. Isolation and production of cells suitable for human therapy: challenges ahead. *Cell Stem Cell*. 2009;4(1):20–26. doi:10.1016/j.stem.2008.11.012
2. Eaker S, Armant M, Brandwein H, et al. Concise review: guidance in developing commercializable autologous/patient-specific cell therapy manufacturing. *Stem Cells Transl Med*. 2013;2(11):871–883. doi:10.5966/sctm.2013-0050
3. Masud S, Torraca V, Meijer AH. Modeling infectious diseases in the context of a developing immune system. *Curr Top Dev Biol*. 2017;124:277–329. doi:10.1016/bs.ctdb.2016.10.006
4. McNiece IK, Wacker KK, Kurtzberg J, Warkentin PI. Standardization, workforce development and advocacy in cell and gene therapies: a summary of the 2020 Regenerative Medicine InterCHANGE. *Cytotherapy*. 2021;23(10):886–893. doi:10.1016/j.jcyt.2021.02.004
5. Osswald A, Hedrich V, Sommergruber W. 3D-3 Tumor models in drug discovery for analysis of immune cell infiltration. *Methods Mol Biol*. 2019;1953:151–162. doi:10.1007/978-1-4939-9145-7_10
6. Pamies D, Bal-Price A, Chesné C, et al. Advanced good cell culture practice for human primary, stem cell-derived and organoid models as well as microphysiological systems. *ALTEX*. 2018;35(3):353–378. doi:10.14573/altex.1710081
7. Pamies D, Leist M, Coecke S, et al. Guidance document on good cell and tissue culture practice 2.0 (GCCP 2.0). *ALTEX*. 2022;39:30–70. doi:10.14573/altex.2111011
8. Pezzulo AA, Starner TD, Scheetz TE, et al. The air-liquid interface and use of primary cell cultures are important to recapitulate the transcriptional profile of *in vivo* airway epithelia. *Am J Physiol Lung Cell Mol Physiol*. 2011;300(1):L25–L31. doi:10.1152/ajplung.00256.2010

PromoCell®



PromoCell GmbH
Sickingenstr. 63/65
69126 Heidelberg
Germany

info@promocell.com
www.promocell.com

USA/Canada

Phone: 1 – 866 – 251 – 2860 (toll free)
Fax: 1 – 866 – 827 – 9219 (toll free)

Deutschland

Telefon: 0800 – 776 66 23 (gebührenfrei)
Fax: 0800 – 100 83 06 (gebührenfrei)

France

Téléphone: 0800 – 90 93 32 (ligne verte)
Téléfax: 0800 – 90 27 36 (ligne verte)

United Kingdom

Phone: 0800 96 03 33 (toll free)
Fax: 0800 169 85 54 (toll free)

Other Countries

Phone: +49 6221 – 649 34 0
Fax: +49 6221 – 649 34 40

© PromoCell GmbH